



Borno - COVID-19 Risk Related Indicators

Assessment of Hard-to-Reach Areas in Northeast Nigeria

May 16-31 2020

Introduction

The continuation of conflict in Northeast Nigeria has created a complex humanitarian crisis, rendering sections of Borno State as hard to reach (H2R) for humanitarian actors. Previous assessments illustrate how the conflict continues to have severe consequences for people in H2R areas. People living in H2R areas who are already facing severe and extreme humanitarian needs are even more vulnerable to the spread of COVID-19, especially due to the lack of health care services and information sources. The first confirmed case in Borno state was announced on 20 April 2020.¹ Due to the limited access to H2R areas it is unlikely there will be confirmation of an outbreak in these areas. It is therefore of utmost importance to evaluate the situation of the population in H2R areas in order to monitor changes and inform humanitarian aid actors on immediate needs of the communities.

Methodology

Using its Area of Knowledge (AoK) methodology, REACH remotely monitors the situation in H2R areas through monthly multi-sector interviews in accessible Local Government Area (LGA) capitals with the following typology of key informants (KIs):

- KIs who are newly arrived internally displaced persons (IDPs) who have left a hard-to-reach settlement in the last 3 months²
- KIs who have had contact with someone living or having been in a hard-to-reach settlement in the last month (traders, migrants, family members, etc.)²

Selected KIs are purposively sampled and are interviewed on settlement-wide circumstances

in H2R areas. Responses from KIs reporting on the same settlement are then aggregated to the settlement level. The most common response provided by the greatest number of KIs is reported for each settlement. When no most common response could be identified, the response is considered as 'no consensus'. While included in the calculations, the percentage of settlements for which no consensus was reached is not always displayed in the results below.

Due to precautions related to the COVID-19 outbreak, data was collected remotely through phone based interviews with assistance from local stakeholders.


Results presented in this factsheet, unless otherwise specified, represent the proportion of settlements assessed within a LGA. Findings are only reported on LGAs where at least 5% of populated settlements and at least 5 settlements in the respective LGA have been assessed. The findings presented are indicative of broader trends in assessed settlements in May 2020, and are not statistically generalisable.³

Assessment Coverage

- 197** Key Informants interviewed
- 108** Settlements assessed
- 10** LGAs assessed
- 5** LGAs with sufficient coverage⁴

Assessment Coverage

Proportion of settlements assessed:



COVID-19 Precautions in IDP Camps

Precautions for New Arrivals

Hand washing and temperature screenings for new arrivals at IDP camps could help slow the spread of COVID-19. To assist in monitoring the implementation of these procedures, REACH began asking KIs, who had recently left H2R areas, if they were asked to wash or sanitise their hands or had their temperature measured when they arrived at the IDP camp.

62% of surveyed KIs had left a H2R area within the last one month, among them:

72% reported they were asked to wash and/or sanitise their hands when they arrived at the IDP camp

23% reported their temperature was measured when they arrived at the IDP camp

¹Nigerian Centre for Disease Control Twitter feed


²Where possible, only KIs that have arrived very recently (0-3 weeks prior to data collection) were interviewed. If not stated otherwise, the recall period is set to one month prior to the last information the KI has had from the hard-to-reach area.

³Due to changes in migration patterns, the specific settlements assessed within each LGA varies each month. Changes in results reported in this factsheet, compared to previous factsheets, may therefore be due to changes in which settlements were assessed instead of changes over time.

⁴LGA level data is only represented for LGAs in which at least 5% of populated settlements and where at least 5 settlements have been assessed. The most recent version of the VTS dataset (released in February 2019 on vts.eocng.org) has been used as the reference for settlement names and locations, and adjusted for deserted villages (OCHA 2020).

Hand Washing Practices in H2R Areas

Proportion of assessed settlements by reported most common hand washing materials by LGA:






Borno - COVID-19 Risk Related Indicators

Assessment of Hard-to-Reach Areas in Northeast Nigeria

May 16-31 2020

Access to Water

Proportion of assessed settlements reporting taking less than 30 minutes to reach their main water source, access water and return to their homes:




Being far from their main water source could potentially mean that households have less access to water and, therefore, less water for hand washing and other hygiene practices. Additionally, travelling to fetch water increases the chance of interacting with others outside of the household along the way to the water point and at the water point, further increasing the risk of transmission.

47% of assessed settlements that reported no access to functional health facility (93%), reported the main reason was that they never existed


Access to Health Care Services

Proportion of assessed settlements reporting a functional health service that the population could reach and return from in one day:



Most Common Health Problems

Proportion of assessed settlements by reported most common health problem, by LGA:⁴



The reported low access to a functional health facility, combined with limited local isolation practices and people's precarious health status, limit their ability to respond to a local COVID-19 outbreak.

COVID-19 Related Symptoms


Proportion of assessed settlements reporting symptoms related to COVID-19, by LGA:

	Breathing difficulties	Coughing	Fever and breathing difficulties	Fever and coughing	None
Bama	11%	11%	11%	11%	84%
Damboa	0%	0%	0%	0%	100%
Gwoza	4%	20%	8%	12%	56%
Konduga	38%	38%	0%	13%	63%
Ngala	0%	0%	0%	0%	100%

Although other viruses and bacteria can cause the three main symptoms associated with COVID-19 (fever, coughing, breathing difficulties), an increase in the reporting of these symptoms could suggest a local COVID-19 outbreak in the H2R areas.

REACH added this indicator to the assessment on 1 April 2020. The proportion of assessed settlements where the symptoms were reported is similar or slightly higher than the proportion reported for each LGA in the 1-15 May factsheet. This is the first time Konduga is being reported on with regards to this indicator.

⁴ These findings represent self-reported health problems. In particular, the fever/malaria responses include fever that is believed to be caused by malaria or other symptoms that the respondents believed are associated with malaria.




Borno - COVID-19 Risk Related Indicators

Assessment of Hard-to-Reach Areas in Northeast Nigeria

May 16-31 2020

Caretaking Practices

Proportion of assessed settlements reporting men/women as primary caretaker when someone is sick, by LGA:




The majority of assessed settlements reported women to be the primary caretaker. This suggests that, based on this indicator, women in the H2R settlements are at higher risk than men of getting infected with COVID-19 in the case of a local COVID-19 outbreak.

22% of assessed settlements reported sick members of the community being separated from others

Communication- Sources of Information

Proportion of assessed settlements by reported main source of information for most people, by LGA:



73% of assessed settlements reported most people had difficulty accessing information they needed regarding available humanitarian assistance

49% of assessed settlements reported community members could listen to the radio

10% of assessed settlements reported community members could use a cell phone


Communication- Information

Communication Access


Communicating COVID-19 related preventative measures, symptoms and when to seek medical care are critical to reduce transmission rates and case fatality ratios. The findings suggest that options to communicate with people in H2R areas are incredibly limited, preventing the dissemination of information and recommendations on COVID-19. However, 56% of the settlements assessed reported that people have heard

about the new coronavirus disease. Similar to the last reporting period, 25% or more of assessed settlements in every LGA reported people have heard about COVID-19. The reported access to radio (49%) and reliance on radio as the main source of information in some areas (overall 23%) suggests that radio campaigns focused on these areas may be able to inform people living in these areas about COVID-19.

Proportion of assessed settlements reporting that most people had received information about the following topics in IDP camps, by LGA:



Proportion of assessed settlements reporting people have heard about COVID-19, by LGA:



Conclusion

The reported lack of access to functional health facilities, existing health problems and limited local isolation practices highlighted in this factsheet, along with the reported lack of use of soap, put communities in H2R areas at a higher risk of infection in the event of a local COVID-19 outbreak. Close monitoring of COVID-19 related symptoms may allow for the prediction of local COVID-19 outbreaks. Additionally, the reported limitations of communication to H2R areas suggest that many people will not have the knowledge they need to prepare for and respond to a local COVID-19 outbreak.